

Main Office 1672-1674 Sheepshead Bay Rd Brooklyn, NY 11235 T- (718) 332 - 0600	Bushwick Office 129 St. Nicholas Ave Brooklyn, NY 11237 T- (718) 821-0643	Endoscopy Center 214 Avenue P (West 6 th & 7 th St) Brooklyn, NY 11204 T- (718) 339 - 5678	Mount Sinai Hospital Brooklyn 3201 Kings Hwy Brooklyn, NY 11234 T- (718) 951 – 2801	Maimonides Medical Center 1025 48 th Street Brooklyn, NY 11219 T- (718) 283-7055	Wyckoff Heights Medical Center 374 Stockholm Street Brooklyn, NY 11237 T- (718) 963-7272	Staten Island Office 2043 Richmond Avenue Staten Island, NY 10314 T- (718) 987-1700	Bronx Office 3594 East Tremont Ave, Suite#100 (LL) Bronx, NY 10465 T- (718) 292 - 3822
---	---	---	---	---	--	---	---

Colonoscopy – CLENPIQ Instructions

APPOINTMENT INFORMATION

Patient Name: _____

Date: _____

Time: _____

Cancellation Policy

Scheduling procedures requires careful planning between the staff, doctors, and facility. Please be compliant with and on-time for your appointment. If you must reschedule or cancel, please contact us at least **1 day prior to procedure**. Failure to do so will incur **A FEE OF \$100**.
Thank you for your cooperation.

1 WEEK PRIOR

- Fill your prescription for CLENPIQ at the pharmacy.
- Arrange for a driver to take you home on the day of the procedure.
- If you have diabetes, check with your doctor regarding diet and medication instructions.
- Avoid all seeds, high fiber vegetables, and peel of all fruits.
- Stop Fiber Supplements, Vitamin E, and Iron containing supplements (example: Ferrous Sulfate)

Medications to be **DISCONTINUED**

- **All Blood thinners** should be discussed with your prescribing doctor. (Coumadin, Plavix, Eliquis, Effient, Brilinta, Xarelto, Arixtra, Lovenox, Pradaxa, Fragmin). You can **CONTINUE Aspirin 81 mg** unless instructed by your doctor.
- **STOP 1 WEEK PRIOR:** Semaglutide (Ozempic, Wegovy), Mounjaro, Zepbound, Trulicity, Victoza, Bydureon BCise
- Fiber supplements and iron supplements
- **STOP 1 DAY PRIOR:** Rybeslsus, Saxenda, Byetta, Adlyxin

1 DAY PRIOR

- Start a clear liquid diet in the morning.
 - **NO SOLID FOOD ALL DAY** (No breakfast, lunch, or dinner)
 - **NO DAIRY PRODUCTS**
 - **NO RED or PURPLE FOOD COLORING**

Clear liquids include: Broths, Water, Clear fruit juices (apple), Soda, Gatorade, Jell-O, Popsicles, Black Coffee or Tea

- At **5PM (PART 1)**: drink 1 bottle of CLENPIQ then drink at least 5 cups (8-ounces each) of clear liquid at your own pace.
- At **9PM (PART 2)**: please repeat the same instruction as (PART 1) with the second bottle of CLENPIQ.

NOTHING TO EAT OR DRINK AFTER MIDNIGHT PRIOR TO THE PROCEDURE.

PROCEDURE DAY

- You may take your morning medications (**High Blood Pressure and Thyroid**) with a small sip of water.
- If you have asthma, please bring your inhaler with you.
 - **NO CHEWING GUM or SUCKING ON HARD CANDY**

A RESPONSIBLE ADULT MUST ACCOMPANY ALL PATIENTS THE DAY OF THE PROCEDURE. NO PATIENT SHOULD DRIVE ON THE DAY OF THE COLONOSCOPY.