

Igor Grosman, DO; F.A.S.G Alexander Brun, MD Prateek Chapalamadugu, MD Nancy Chen, MD

Konstantin Peysin, DO

Yevgeniy Mikityanskiy, DO

Dmitriy Khodorskiy, MD

Board Certified in Gastroenterology & Hepatology

E – Mail: info@triboroughgi.com, www.triboroughgi.com

Main Office 1672-1674 Sheepshead Bay Rd Brooklyn, NY 11235 T- (718) 332 - 0600

**Bushwick Office** 129 St. Nicholas Ave Brooklyn, NY 11237 T- (718) 821-0643

Patient Name:

 e
 Endoscopy Center

 Ave
 214 Avenue P

 37
 (West 6<sup>th</sup> & 7<sup>th</sup> St)

 Brooklyn, NY 11204
 T- (718) 339 - 5678

Mount Sinai Hospital Brooklyn 3201 Kings Hwy Brooklyn, NY 11234 T- (718) 951 – 2801 Maimonides Medical Center 1025 48<sup>th</sup> Street Brooklyn, NY 11219 T- (718) 283-7055 Wyckoff Heights Medical Center 374 Stockholm Street Brooklyn, NY 11237 T- (718) 963-7272

 s
 Staten Island Office

 2043 Richmond Avenue
 2043 Richmond Avenue

 staten Island, NY 10314
 37

 T- (718) 987-1700
 2

Bronx Office 3594 East Tremont Ave, Suite#100 (LL) Bronx, NY 10465 T- (718) 292 - 3822

# **Colonoscopy – PLENVU Instructions**

APPOINTMENT INFORMATION

# **Cancellation Policy**

Scheduling procedures requires careful planning between the staff, doctors, and facility. Please be compliant with and on-time for your appointment. If you must reschedule or cancel, please contact us at least **1 day prior to procedure**. Failure to do so will incur **A FEE OF \$100**. Thank you for your cooperation.

# **1 WEEK PRIOR**

Date:

Time: \_\_\_\_

- Fill your prescription for PLENVU at the pharmacy.
- Arrange for a driver to take you home on the day of the procedure.
- If you have diabetes, check with your doctor regarding diet and medication instructions.
- Avoid all seeds, popcorn, multigrain breads, high fiber vegetables, and fruits.
- Stop Fiber Supplements, Vitamin E, and Iron containing supplements (example: Ferrous Sulfate)

# Medications to be **DISCONTINUED**

- All Blood thinners should be discussed with your prescribing doctor. (Coumadin, Plavix, Eliquis, Effient, Brilinta, Xarelto, Arixtra, Lovenox, Pradaxa, Fragmin). You can CONTINUE Aspirin 81 mg unless instructed by your doctor.
- STOP 1 WEEK PRIOR: Semaglutide (Ozempic, Wegovy), Mounjaro, Zepbound, Trulicity, Victoza, Bydureon BCise
- o STOP 1 DAY PRIOR: Rybeslsus, Saxenda, Byetta, Adlyxin

# **1 DAY PRIOR**

- Start a clear liquid diet in the morning.
  - **NO SOLID FOOD ALL DAY** (No breakfast, lunch, or dinner)
  - **o** NO DAIRY PRODUCTS
  - NO RED or PURPLE FOOD COLORING

Clear liquids include: Broths, Water, Clear fruit juices (apple), Soda, Gatorade, Jell-O, Popsicles, Black Coffee or Tea

- At <u>5PM</u> (**PART 1**): Add Dose 1 (pouch 1) into the mixing container then fill the container up to the 16-ounces line with water. DRINK ALL THE LIQUID in the container within 30 minutes. In addition, drink 16-ounces of water over the next 1 hour.
- At <u>9PM</u> (PART 2): Add Dose 2 (pouch A and pouch B) into the mixing container and repeat the same instruction as (PART 1) and drink clear liquids until midnight.

### NOTHING TO EAT OR DRINK AFTER MIDNIGHT PRIOR TO THE PROCEDURE.

### **PROCEDURE DAY**

- You may take your morning medications (High Blood Pressure and Thyroid) with a small sip of water.
- If you have asthma, please bring your inhaler with you.
  - NO CHEWING GUM or SUCKING ON HARD CANDY

A RESPONSIBLE ADULT MUST ACCOMPANY ALL PATIENTS THE DAY OF THE PROCEDURE. NO PATIENT SHOULD DRIVE ON THE DAY OF THE COLONOSCOPY.